

THE COLOGUARD® COLLECTION KIT

The Cologuard Collection Kit is easy to use. If a patient has questions, a live Customer Support Specialist is available to assist via phone: **1-844-870-8870**.

What's inside:



SHIPPING BOX



SAMPLE CONTAINER



PROBE + TUBE



BRACKET



PATIENT GUIDE INSTRUCTIONS

Step 6: Label Your Samples

You **must** label your samples with the information listed below. Use a ballpoint pen and write on a hard, flat surface.

EXAMPLE

First Name	First Name	First Name
Last Name	Last Name	Last Name
DOB (MM/DD/YY)	DOB (MM/DD/YY)	DOB (MM/DD/YY)
Date of Birth (mm/dd/yyyy)	Date of Birth (mm/dd/yyyy)	Date of Birth (mm/dd/yyyy)
Date of Collection (mm/dd/yyyy)	Date of Collection (mm/dd/yyyy)	Date of Collection (mm/dd/yyyy)
Site 1 (1-3)	Site 1 (1-3)	Site 1 (1-3)
Time of Collection (00:00 AM/PM)	Time of Collection (00:00 AM/PM)	Time of Collection (00:00 AM/PM)

a. Fill out both labels. **Print** your information as shown, in the example.

b. Peel the label away from the paper.

c. Wrap one label around the probe tube as shown.

d. Place one label on the lid of the stool sample container as shown.

For questions or help call Exact Sciences at 1-844-870-8870
www.cologuard.com
Rx only

Page 1 of 1

SAMPLE LABELS



BOTTLE OF LIQUID
(PRESERVATIVE)

Patients can watch a short, helpful video on how to use the collection kit at [Cologuard.com/use](https://www.cologuard.com/use).

EASY STEPS FOR SAMPLE COLLECTION

Patients collect a single stool sample at home and send to Exact Sciences Labs via prepaid UPS shipping or pick-up

1 SIT



Cologuard is ready to use when you are ready to use it.

2 SCRAPE



Use the kit components to collect your sample.

3 SOAK



Fill the sample container with the liquid preservative.

4 SHIP



Ship your sample to complete the process.

Indications and Important Risk Information

Cologuard is intended for the qualitative detection of colorectal neoplasia associated DNA markers and for the presence of occult hemoglobin in human stool. A positive result may indicate the presence of colorectal cancer (CRC) or advanced adenoma (AA) and should be followed by diagnostic colonoscopy. Cologuard is indicated to screen adults of either sex, 45 years or older, who are at typical average risk for CRC. Cologuard is not a replacement for diagnostic colonoscopy or surveillance colonoscopy in high-risk individuals.

Cologuard is not for high-risk individuals, including patients with a personal history of colorectal cancer and adenomas; have had a positive result from another colorectal cancer screening method within the last 6 months; have been diagnosed with a condition associated with high risk for colorectal cancer such as IBD, chronic ulcerative colitis, Crohn's disease; or have a family history of colorectal cancer, or certain hereditary syndromes.

Positive Cologuard results should be referred to diagnostic colonoscopy. A negative Cologuard test result does not guarantee absence of cancer or advanced adenoma. Following a negative result, patients should continue participating in a screening program at an interval and with a method appropriate for the individual patient.

False positives and false negatives do occur. In a clinical study, 13% of patients without colorectal cancer or advanced adenomas received a positive result (false positive) and 8% of patients with cancer received a negative result (false negative). The clinical validation study was conducted in patients 50 years of age and older. Cologuard performance in patients ages 45 to 49 years was estimated by sub-group analysis of near-age groups.

Cologuard performance when used for repeat testing has not been evaluated or established. Rx only.

Order forms are available online at CologuardHCP.com/resources or by calling **1-844-870-8870**.